

AD-A125 044

ROBUSTNESS PROPERTIES OF THE F-TEST AND BEST LINEAR
UNBIASED ESTIMATORS I. (U) PITTSBURGH UNIV PA CENTER
FOR MULTIVARIATE ANALYSIS B K SINHA ET AL. NOV 82

1/1

UNCLASSIFIED


TR-82-28 AFOSR-TR-83-0018 F49620-82-K-0001 F/G 12/1

NL

END


FILMED

DTIC




MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A125044



Center for Multivariate Analysis
University of Pittsburgh



DTIC
ELECTE
MAR 01 1983
E

DTIC FILE COPY

83 02 028 151

Approved for public release;
distribution unlimited

①

**ROBUSTNESS PROPERTIES OF THE \hat{F} -TEST AND
BEST LINEAR UNBIASED ESTIMATORS
IN LINEAR MODELS**

Bimal Kumar Sinha¹

and


Hilmar Drygas²

University of Pittsburgh

November 1982

Technical Report No. 82-28

Center for Multivariate Analysis
University of Pittsburgh
Ninth Floor, Schenley Hall
Pittsburgh, PA 15260




¹Work of this author is sponsored by the Air Force Office of Scientific Research under Contract F49620-82-K-0001. Reproduction in whole or in part is permitted for any purpose of the United States Government.

²On leave from Fachbereich 17 (Mathematik) der Gesamthochschule, Postfach 101 380, D-3500 Kassel, F.R. Germany.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC

This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.

MATTHEW J. KERPER
Chief, Technical Information Division



Accession For

GRA&I

TAB

nounced

Justification

By
Distribution/

Availability Code

Dist Avail and/or
Special

A

ROBUSTNESS PROPERTIES OF THE F-TEST AND BEST LINEAR
UNBIASED ESTIMATORS IN LINEAR MODELS

by

Bimal Kumar Sinha^{1,3}

and

Hilmar Drygas^{1,2} *epsilon prime*

ABSTRACT

Consider a linear model $Y = X\beta + \epsilon$, $E(\epsilon) = 0$, $E(\epsilon\epsilon') = I_n$ with β unknown. For the problem of testing the linear hypothesis $C\beta = 0$, $\text{im}(C') \subseteq \text{im}(X')$, Ghosh and Sinha (1980) proved that the properties of the usual F-test being LRT and UMPI (under a suitable group of transformations) remain valid for specific non-normal families. In this paper it is shown that both criterion and inference robustness of the F-test hold under the assumption $\epsilon \sim q(\epsilon)$, q convex and isotonic. This result is similar to a robustness property of Hotelling's T^2 -test proved by Kariya (1981). Finally it is proved that the Best Linear Unbiased Estimator (BLUE) of any estimable function $C\beta$ is more concentrated around $C\beta$ than any other unbiased estimator of $C\beta$ under the assumption that ϵ is spherically distributed.

Key words: Linear model, linear hypothesis, spherical distribution, estimable function, LRT, UMPI, BLUE.

¹Department of Mathematics and Statistics, University of Pittsburgh, Pittsburgh, PA 15260, U.S.A.

²On leave from Fachbereich 17 (Mathematik) der Gesamthochschule, Postfach 101 380, D-3500 Kassel, F.R. Germany.

³Work of this author is sponsored by the Air Force Office of Scientific Research under Contract F49629-82-K-001. Reproduction in whole or in part is permitted for any purpose of the United States Government.

1. INTRODUCTION

We consider the linear model

$$Y = X\beta + \sigma\epsilon, \quad (1.1)$$

where Y is $n \times 1$, X is $n \times k$, β is $k \times 1$ and $\sigma > 0$. It is assumed that $\text{Rank}(X) = r \leq k < n$ and that

$$E\epsilon = 0, \text{Cov } \epsilon = E(\epsilon\epsilon') = I_n. \quad (1.2)$$

Then

$$EY = X\beta, \text{Cov } Y = \sigma^2 I_n. \quad (1.3)$$

We denote by $\text{im}(B)$ the column-space of a matrix B .

We want to consider linear hypotheses of the kind $H_0: C\beta = \delta$ where C is a $s \times k$ -matrix, $\text{Rank}(C) = s$. In order that these hypotheses are testable it is necessary that $\text{im}(C') \subseteq \text{im}(X')$, i.e., $C = TX$ for some T . It is well-known that the standard F -test is both LRT (Likelihood-Ratio-Test) and UMPI (uniformly most powerful invariant) under the normality of the vector ϵ with respect to a suitable group of affine transformations. Ghosh and Sinha (1980) proved that both of these properties of the F -test remain valid even under typical non-normal distributions of the errors, namely normal (scale) mixtures which include multivariate t -distributions. The purpose of this paper is to prove an even stronger result. It is shown that the F -test is both LRT and UMPI under the distributional assumption

$$\varepsilon \sim q(\varepsilon' \varepsilon), \quad q \uparrow, \text{ convex} \quad (1.4)$$

(\uparrow describes an isotonic function). The results are similar to a robustness property proved for Hotelling's T^2 -test by Kariya (1981).

In section 2 we present a canonical form of the problem and section 3 deals with the UMPI property of the F-test. To obtain the distribution of a maximal invariant, certain results due to Dawid (1977), (see also Dempster (1969)) and Wijsman's representation theorem (Wijsman, 1967) are used. The property of the F-test being LRT and the distribution of the F-statistics under (1.4) are indicated in section 4. Finally, in section 5 it is shown for a more general linear model with spherically distributed error that the BLUE (Best Linear Unbiased Estimator) of an estimable function $C\beta$ is more concentrated around $C\beta$ than any other unbiased estimator (robustness property of the BLUE).

2. A CANONICAL FORM

According to our assumption $\text{rank}(X) = r \leq k < n$. Then there exists (see, e.g., Rao (1973)) a quasi-orthogonal matrix P_1 of order $k \times n$ such that $P_1 P_1' = I_k$, and $k \times k$ -diagonal matrix Λ and an orthogonal $k \times k$ -matrix Q such that the singular value decomposition

$$X = P_1' \Lambda Q \quad (2.1)$$

holds. Here $\Lambda^2 = \text{diag}(\lambda_1^2, \dots, \lambda_r^2, 0, \dots, 0)$ and the λ_i^2 are the positive eigenvalues of $X'X$. Supplement P_1 by P_2 , a matrix of

order $n - k \times n$ such that $P = (P_1' : P_2')'$ is an orthogonal $n \times n$ -matrix.

Let us now consider the transformation

$$Y \rightarrow \begin{cases} Y_{(1)} = P_1 Y = P_1 X \beta + \sigma P_1 \epsilon = \Lambda \gamma + \sigma \epsilon_1^* \\ Y_{(2)} = P_2 Y = \sigma P_2 \epsilon = \sigma \epsilon_1^* \end{cases} \quad (2.2)$$

where

$$\epsilon_1^* = P_1 \epsilon, \quad \epsilon_2^* = P_2 \epsilon, \quad \gamma = Q \beta = (\gamma_1, \dots, \gamma_k)'. \quad (2.3)$$

Note that $\Lambda \gamma = \text{diag}(\lambda_1 \gamma_1, \dots, \lambda_r \gamma_r, 0, \dots, 0)$. We denote the $n-r$ components of the vector $Y = (Y_{(1)}', Y_{(2)}')'$ which have mean zero by u_1, \dots, u_{n-r} . The vector of the first r components of $Y_{(1)}'$ which have possibly a non-zero mean, is denoted by $Y_{(11)}'$. Since $C = TX$, the equation $C \beta = \delta$ is equivalent to $TX \beta = TP_1' \Lambda Q \beta = TP_1' \Lambda \gamma = \delta$. Now let us split up P_1 as $P_1 = (P_{11}, P_{12})$, where P_{11} is $n \times r$ and P_{12} is $n \times k-r$. Let $T_1 = T P_{11}'$. Then clearly $\text{rank}(T_1) = s$ and $TX \beta = \delta$ is equivalent to $T_1 \text{diag}(\lambda_1 \gamma_1, \dots, \lambda_r \gamma_r) = \delta$. Let T_2 be a $r-s \times r$ -matrix such that $\text{Rank}(T_2) = r-s$ and $T_2' T_1 = 0$. We now employ the following transformation:

$$Y_{(11)} \rightarrow \begin{cases} (T_1 T_1')^{-1} (T_1 Y_{(11)}) - \theta = (v_1, \dots, v_s)' \\ (T_2 T_2')^{-1} (T_2 Y_{(11)}) = (w_1, \dots, w_{r-s})' \end{cases} \quad (2.4)$$

Combining (2.2) and (2.4) we can conclude the following:

$$\begin{aligned} v &= (v_1, \dots, v_s)' = \theta + \sigma \epsilon_1 \\ w &= (w_1, \dots, w_{r-s})' = \eta + \sigma \epsilon_2 \\ u &= (u_1, \dots, u_{n-r})' = \sigma \epsilon_3 \end{aligned} \quad (2.5)$$

with $\epsilon = (\epsilon_1', \epsilon_2', \epsilon_3')'$, $E \epsilon = 0$, $E \epsilon \epsilon' = I_n$, $\epsilon \sim q(\epsilon' \epsilon)$, q convex, \dagger and $H_0: A \beta = \delta$ is equivalent to $H_0: \theta = 0$, η and σ^2 being nuisance para-

meters. This is the canonical form of the problem. It may be noted that Ghosh and Sinha (1980) made use of this canonical form without explicitly stating how it can be obtained. The above analysis spells out clearly the necessary steps leading from (1.1) to (2.3). Under the assumption of normality of ε it follows clearly that the F-test based on $F = (v'v)/(u'u)$ is LRT and UMPI (under a suitable group of transformations).

3. UMPI - PROPERTY OF THE F-TEST

Without any loss of generality we treat the problem in the canonical form (2.5). As in Ghosh and Sinha (1980) obviously the testing problem remains invariant under the group G of transformations

$$v \rightarrow c P_s v, w \rightarrow w+b, u \rightarrow c P_{n-r} u, \quad (3.1)$$

where c runs over the non-zero reals, b runs over \mathbb{R}^{r-s} , and P_k runs over p_k , the group of $k \times k$ orthogonal transformations ($k=s$ and $n-r$, respectively). As is well-known a maximal invariant under G is $T_0 = (v'v)/(u'u)$ and a maximal invariant parameter is $\delta = (\theta'\theta)/\sigma^2$. To obtain a formal expression of the distribution function of T_0 , we first note that the joint distribution of v and u has the density (Dawid (1977), Dempster (1969))

$$f(v, u | \theta, \sigma^2) = \sigma^{-(n-r+s)} q([(v-\theta)'(v-\theta) + u'u]/\sigma^2), \quad q \text{ convex, } \hat{1}. \quad (3.2)$$

To obtain the distribution of T_0 from (3.2) we apply Wijsman's (1967) representation theorem which we state below as a lemma in a suitable form:

3.1 Lemma. Let $f(z)$ be the pdf of $z = (v', u')'$ and let $T_0 = t(z)$ be a maximal invariant under the group G of transformations (3.1): $v \rightarrow c P_s v$, $u \rightarrow c P_{n-r} u$. Let, moreover, $P_\delta^{T_0}$ be the distribution induced by T_0 under $\delta = (\theta' \theta) / \sigma^2$. Then the pdf of T with respect to $P_0^{T_0}$ evaluated at $T_0 = t(z)$ is given by

$$\frac{dP_\delta^{T_0}}{dP_0^{T_0}}(t(z)) = f_{T_0}(t(z) | \delta) \quad (3.3)$$

$$= \left(\int_G f(gz | \theta, \sigma^2) |c|^{n-r-s} dv(g) \right) / \left(\int_G f(gz | 0, \sigma^2) |c|^{n-r-s} dv(g) \right)$$

where ν is a left invariant measure on G and $g = (c, P_s, P_{n-r})$, $gz = (c P_s v, c P_{n-r} u)$.

Conditions under which (3.3) is valid are stated in Wijsman (1967). It is easy to verify that these conditions are satisfied in our case [see Kariya (1978, 1981)]. We choose $\nu = \nu_1 \times \nu_2 \times \nu_3$ where $d\nu_1(c) = dc / |c|$ and ν_2 and ν_3 are the left invariant probability measures on P_s and P_{n-r} , respectively. Lemma 3.1 applied to our setup (3.2) yields the following.

3.2 Lemma. The pdf of T_0 , $f_{T_0}(t | \delta)$ is evaluated as

$$\left(\int_{-\infty}^{\infty} \int_{P_s} q(c^2 - 2c[(v' P_s \eta) / (v' v + \eta' \eta)]^{1/2} + \eta' \eta) dv_2(P_s) |c|^{n-r+s-1} dc \right) / \left(\int_{-\infty}^{\infty} q(c^2) |c|^{n-r+s-1} dc \right), \quad (3.4)$$

where $\eta = \theta / \sigma$ and $\eta' \eta = \delta$.

Proof. From (3.2) and the action of the transformation g on $z = (v', u')'$ we get

$$\begin{aligned}
f(g \cdot z | \theta, \sigma^2) &= \sigma^{-(n-r+s)} q([(c P_S v - \theta)' (c P_S v - \theta) + c^2 u' u] / \sigma^2) \\
&= \sigma^{-(n-r+s)} q([c^2 (v' v + u' u) - 2 c v' P_S' \theta + \theta' \theta] / \sigma^2) \\
&= \sigma^{-(n-r+s)} q((c^*)^2 - 2 c^* (v' v + u' u)^{-\frac{1}{2}} v' P_S' \eta + \eta' \eta)
\end{aligned}$$

where $c^* = [c(v' v + u' u)^{\frac{1}{2}}] / \sigma$.

From (3.3) the numerator of $f_{T_0}(t | \delta)$ is obtained as

$$\int_{-\infty}^{\infty} \int_{P_{r-s}} \int_{P_S} q((c^*)^2 - 2 c^* v' P_S' \eta (v' v + u' u)^{-\frac{1}{2}} + \eta' \eta) |c^*|^{n-r+s-1} (v' v + u' u)^{-(n-r+s)/2} dc^* dv_2(P_S) dv_3(P_{r-s}) \quad (3.5)$$

whereas the denominator of (3.3) obtained from (3.5) by putting $\eta = 0$, is given by

$$\int_{-\infty}^{\infty} \int_{P_{r-s}} \int_{P_S} q((c^*)^2) |c^*|^{n-r+s-1} (v' v + u' u)^{-(n-r+s)/2} dc^* dv_2(P_S) dv_3(P_{r-s}) \quad (3.6)$$

Since the integrand in (3.5) does not depend on P_{r-s} and the integrand in (3.6) does not depend on P_S and P_{r-s} and the integral $\int_{r-s} dv_3(P_{r-s})$ equals one, together with $\int_{P_S} dv_2(P_S) = 1$, by taking the ratio of (3.5) to (3.6), the lemma follows. Q.E.D.

We are now in a position to prove our main result.

3.3 Theorem. Under the model (2.3), the F-test based on T_0 is UMPI with respect to the group G of transformations (3.1).

Proof. Using the convexity of q we will show that $f_{T_0}(t | \delta)$ for fixed $\delta > 0$ is isotonic in t ($\uparrow t$), thereby proving the theorem.

Let $t^* = v(v' v + u' u)^{-\frac{1}{2}}$ implying that $f_{T_0}(t | \delta)$ evidently depends on t^* . We first show that $f_{T_0}(t | \delta) \Big|_{t^*} = f_{T_0}(t | \delta) \Big|_{0(s)t^*}$ for

all orthogonal matrices $O(s) \in P_s$. This will mean that f really depends on $(t^*)'t^*$, i.e., on t . Replacing t^* by $O(s)t^*$ we get the numerator of $f_{T_0}(t|\delta)$ in (3.4) as

$$\int_{-\infty}^{\infty} \int_{P_s} q(c^2 - 2c v'(O(s))' P_s \eta (v'v + u'u)^{-\frac{1}{2}} + \eta'\eta) |c|^{n-r+s-1} dv_2(P_s) dc.$$

Since $(O(s))'P_s$ is orthogonal and $v_2(P_s) = v_2((O(s))'P_s)$ due to the left-invariance property of the measure v the claim follows.

Denoting the numerator of $f_{T_0}(t|\delta)$ by $H(t^*|\delta)$, then in view of $H(t^*|\delta) = H(-t^*|\delta)$ it follows from the convexity of q that for any $\alpha \in [0,1]$:

$$\begin{aligned} H(-t^*|\delta) &= H(t^*|\delta) = \alpha H(t^*|\delta) + (1-\alpha) H(-t^*|\delta) \\ &= \int_{-\infty}^{\infty} \int_{P_s} [\alpha q(c^2 - 2c(t^*)' P_s \eta + \eta'\eta) \\ &\quad + (1-\alpha) q(c^2 + 2c((t^*)' P_s \eta + \eta'\eta))] |c|^{n-r+s-1} dv_2(P_s) dc \\ &\geq \int_{-\infty}^{\infty} \int_{P_s} [q(c^2 - 2c(2\alpha-1)(t^*)' P_s \eta + \eta'\eta)] |c|^{n-r+s-1} dv_2(P_s) dc \\ &= H((2\alpha-1)t^*|\delta) = H(-(2\alpha-1)t^*|\delta). \end{aligned} \tag{3.7}$$

Since $H(t^*|\delta)$ depends only on $(t^*)'t^*$ and $(2\alpha-1)^2 \leq 1$ it follows indeed that $f_{T_0}(t|\delta)$ is isotonic. Q.E.D.

4. LRT - PROPERTY OF THE F-TEST AND ITS DISTRIBUTION

Since $q \uparrow$, it follows that the maximum likelihood estimators of θ, η are given by $\hat{\theta} = v, \hat{\eta} = w$, yielding

$$\sup_{\theta, \eta} L(\theta, \eta, \sigma^2 | v, w, u) = \sigma^{-n} q([u'u]/\sigma^2). \quad (4.1)$$

In order to maximize the right hand side of (4.1) with respect to σ it is easily seen that

$$\sup_{\sigma} \sigma^{-n} q([u'u]/\sigma^2) = (u'u)^{-n/2} K \quad (4.2)$$

where $K = \sup_{\sigma} \sigma^{-n} q(\sigma^{-2})$.

Similarly the supremum of the likelihood function under $H_0: \theta = 0$ is obtained as

$$\sup_{\eta, \sigma} \sigma^{-n} q([v'v + (w-\eta)'(w-\eta)]/\sigma^2) = (u'u + v'v)^{-n/2} K. \quad (4.3)$$

Hence the LRT is equivalent to the F-test.

That the F-statistic $F = (v'v)(n-r)/(u'u)s$ is distributed as a central F-variable under H_0 whatever q may be follows from the results due to Dawid (1977).

5. A ROBUSTNESS PROPERTY OF THE BEST LINEAR UNBIASED ESTIMATOR

Consider the linear model

$$y = X\beta + U\epsilon, \quad E(\epsilon) = 0, \quad E(\epsilon\epsilon') = I_m, \quad (5.1)$$

where X is a given $n \times k$ and U is a given $n \times m$ -matrix, $\text{Rank}(U) = m$. Then

$$E y = X\beta, \quad \text{Cov } y = E(y - E y)(y - E y)' = U U' =: Q. \quad (5.2)$$

Let $C\beta$ be any estimable function of β , i.e., $C = TX$ for some T .

A best linear unbiased estimator of $C\beta$ is any linear function G such that $GX = C$ and $GQa = 0$ if $X'a = 0$.

In the sequel we will assume that ϵ is spherically distributed. This means that $T\epsilon$ has the same distribution as ϵ for all orthogonal $m \times m$ matrices T . If ϵ has a density this is equivalent to the requirement that the density is a function of $\epsilon'\epsilon$. This follows from the fact that $\epsilon'\epsilon$ is a maximal invariant with respect to the group of orthogonal transformations.

We will show that any BLUE Gy of $C\beta$ is more concentrated around $C\beta$ than any other unbiased estimator Ly of $C\beta$. More precisely:

5.1 Theorem. Let the model (5.1) be given and let ϵ be spherically distributed. Then for any $c \in \mathbb{R}$ and any n.n.d. matrix A of appropriate order

$$\begin{aligned} P((Gy - C\beta)' A (Gy - C\beta) \leq c^2) &\geq \\ P((Ly - C\beta)' A (Ly - C\beta) \leq c^2), \end{aligned} \quad (5.3)$$

where Gy is any BLUE of $C\beta$ and Ly is any unbiased estimator of $C\beta$.

Proof. For any unbiased estimator Ly of $C\beta$ we get $Ly = LX\beta + LU\epsilon = C\beta + LU\epsilon$. Therefore the two probabilities which have to be compared are:

$$P((\epsilon'U'G'AGU\epsilon) \leq c^2) \text{ and } P((\epsilon'U'L'ALU\epsilon) \leq c^2). \quad (5.4)$$

We prove that for any L_0 the non-zero eigenvalues of $U'L_0'AL_0U$ and $A^{\frac{1}{2}}L_0UU'L_0'A^{\frac{1}{2}} = A^{\frac{1}{2}}L_0QL_0'A^{\frac{1}{2}}$ coincide. Indeed if $U'L_0'AL_0Ux = \lambda x$, $x \neq 0$, then $\lambda A^{\frac{1}{2}}L_0Ux = A^{\frac{1}{2}}L_0UU'L_0'A^{\frac{1}{2}}L_0Ux = A^{\frac{1}{2}}L_0QL_0'A^{\frac{1}{2}}(A^{\frac{1}{2}}L_0Ux)$ implying that λ is an eigenvalue of $A^{\frac{1}{2}}L_0QL_0'A^{\frac{1}{2}}$ unless $A^{\frac{1}{2}}L_0Ux = 0$

in which case $\lambda = 0$ in view of $x \neq 0$. In the same way it follows from $A^{\frac{1}{2}} L_0 Q L_0' A^{\frac{1}{2}} y = \mu y$ by premultiplying with $U' L_0' A^{\frac{1}{2}}$ that μ is an eigenvalue of $U' L_0' A L_0 U$ unless $U' L_0' A^{\frac{1}{2}} y = 0$ in which case again $\mu = 0$.

Since $A^{\frac{1}{2}} G Q G' A^{\frac{1}{2}} \leq A^{\frac{1}{2}} L Q L' A^{\frac{1}{2}}$ by the BLUE-property it follows from the Courant-Fisher min-max-Theorem (see, e.g., Bellman (1970), p. 115-117) that all the eigenvalues of $A^{\frac{1}{2}} G Q G' A^{\frac{1}{2}}$ are smaller than the corresponding eigenvalues of $A^{\frac{1}{2}} L Q L' A^{\frac{1}{2}}$ if ordered according to decreasing magnitude. This would also hold for $U' G' A G U$ and $U' L' A L U$ if we can show that the dimension of the null-space of $U' L' A U$ cannot be greater than the dimension of the null-space of $U' G' A G U$. Now $U' G' A U x = 0$ is equivalent to $A G U x = 0$. This can happen either if $x = U' b$, $X' b = 0$ or $U x = X \beta$ and $A C \beta = A T X \beta = 0$. If $U x = X \beta$ and $T X \beta = C \beta \in \text{im}(A)$ does not vanish unless $C \beta = 0$. But in this case $A L U x = A L X \beta = C \beta$ does not vanish, too. Therefore $\dim(\text{im}(A L U)) \geq \dim(U^{-1}(\text{im} X) \cap (T U)^{-1}(\text{im} A)) = \dim(\text{im}(U' G' A G U))$. This implies the required condition about the null-space. Now let T be an orthogonal transformation such that $U' G' A G U = T \text{diag}(\lambda_1, \dots, \lambda_n) T'$ and similarly T_1 be an orthogonal transformation such that $T_1 \text{diag}(\lambda_1^*, \dots, \lambda_m^*) T_1' = U' L' A L U$. By spherical symmetry $T' \epsilon$ and $T_1' \epsilon$ have the same distribution as ϵ . Therefore

$$P(\epsilon' U' G' A G U \leq c^2) = P(\epsilon' \text{diag}(\lambda_1, \dots, \lambda_m) \epsilon \leq c^2) \quad (5.5)$$

and

$$P(\epsilon' U' L' A L U \epsilon \leq c^2) = P(\epsilon' \text{diag}(\lambda_1^*, \dots, \lambda_m^*) \epsilon \leq c^2). \quad (5.6)$$

Since $\lambda_1 \leq \lambda_1^*$, $i=1,2,\dots,m$, it follows that $\epsilon' \text{diag}(\lambda_1, \dots, \lambda_m) \epsilon \leq \epsilon' \text{diag}(\lambda_1^*, \dots, \lambda_m^*) \epsilon$ and from this evidently (5.3) is obtained.

Q.E.D.

5.2 Remark. A weak version of this result was recently presented by Ali and Ponnappalli, (1982).

REFERENCES

- ALI, MIR M. and PONNAPALLI, R. (1982). "An Optimum Property of Gauss-Markov Estimates when Errors have elliptic distribution". Abstract, IMS Bulletin (July 1982), Paper presented at the Annual Meeting of the American Statistical Association, Cincinnati, 1982.
- BELLMAN, R. (1970). Introduction to Matrix Analysis. 2nd. ed., McGraw Hill Book Company, New York.
- DAWID, A.P. (1977). "Spherical Matrix Distributions and a Multivariate Model". J. Roy. Statist. Soc., Ser. B., 39, 254-261.
- DEMPSTER, A.P. (1969). Elements of Continuous Multivariate Analysis, Addison-Wesley, Reading, Mass.
- GHOSH, M. and SINHA, B.K. (1980). "On the Robustness of Least Squares Procedures in Regression Models". Journal of Multivariate Analysis 10, 332-342.
- KARIYA, T. (1981). "A Robustness property of Hotelling's T^2 -Test". Ann. of Statist. 9, 211-214.
- KARIYA, T. (1978). "The General Manova Problem". Ann. Statist. 6, 200-214.
- LEHMANN, E.L. (1959). Testing Statistical Hypotheses. Wiley, New York.
- RAO, C.R. (1973). Linear Statistical Inference and its Applications. Wiley, New York.
- WIJSMAN, R.A. (1967). "Cross Section of Orbits and their Application to Densities of Maximal Invariants". Fifth Berkeley Symp. Math. Statist. Prob. 1, 389-400.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 83 - 0018	2. GOVT ACCESSION NO. AD-A125 044	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Robustness Properties of the F-Test and Best Linear Unbiased Estimators in Linear Models		5. TYPE OF REPORT & PERIOD COVERED Technical -
7. AUTHOR(s) Bimal Kumar Sinha and Hilmar Drygas		6. PERFORMING ORG. REPORT NUMBER TR- 82-28
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Multivariate Analysis University of Pittsburgh Pittsburgh, PA 15260		8. CONTRACT OR GRANT NUMBER(s) F49620-82-K-0001
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research Department of the Air Force Rolling Air Force Base, DC 20332		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2304/AS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 1982
		13. NUMBER OF PAGES 11
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Lines model, linear hypothesis, spherical distribution, estimable function, LRT, UMPI, BLUE.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Consider a linear model $Y = X\beta + \epsilon$, $E(\epsilon) = 0$, $E(\epsilon\epsilon') = I_n$ with β, σ unknown. For the problem of testing the linear hypothesis $C\beta = \delta$, $\text{im}(C') \subseteq \text{im}(X')$, Ghosh and Sinha (1980) proved that the properties of the usual F-test being LRT and UMPI (under a suitable group of transformations) remain valid for specific non-normal families. In this paper it is shown that both criterion and inference robustness of the F-test hold under the assumption $\epsilon \sim q(\epsilon'\epsilon)$, q convex and isotonic.		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

This result is similar to a robustness property of Hotelling's T^2 -test proved by Kariya (1981). Finally it is proved that the Best Linear Unbiased Estimator (BLUE) of any estimable function $C\beta$ is more concentrated around $C\beta$ than any other unbiased estimator of $C\beta$ under the assumption that ϵ is spherically distributed.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

3-8

DT